

CD24

Concentrated Monoclonal Antibody
Control Number: 901-323-011615

ISO
9001&13485
CERTIFIED

Catalog Number: CM 323 A
Description: 0.1 ml, concentrated
Dilution: 1:100-1:200
Diluent: Van Gogh Yellow

Intended Use:

For In Vitro Diagnostic Use

CD24 [SN3b] is a mouse monoclonal antibody that is intended for laboratory use in the qualitative identification of CD24 protein by immunohistochemistry (IHC) in formalin-fixed paraffin-embedded (FFPE) human tissues. The clinical interpretation of any staining or its absence should be complemented by morphological studies using proper controls and should be evaluated within the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Summary and Explanation:

CD24 (30-70 kDa molecular weight) is a two-chain glycosylphosphatidylinositol (GPI) anchored glycoprotein, that acts as a mucin-like adhesion molecule. CD24 has been shown to functionally enhance the metastatic potential of malignant-cells, as it has been also identified as a ligand of P-selectin, an adhesion receptor on activated endothelial cells and platelets (2). CD24 expression has been reported to be a potentially significant marker for diagnosis of a wide variety of human cancers (3). According to studies, elevated CD24 membranous expression and, in particular, cytoplasmic staining seem to predict malignant transformation. CD24 has been reported as a prognostic marker for breast cancer and more specifically for tamoxifen-resistant breast cancer cases (1). Further studies have shown a subpopulation (CD44+/CD24-) of breast cancer cells to have stem/progenitor cell properties (3).

Principle of Procedure:

Antigen detection in tissues and cells is a multi-step immunohistochemical process. The initial step binds the primary antibody to its specific epitope. After labeling the antigen with a primary antibody, an enzyme labeled polymer is added to bind to the primary antibody. The detection of the bound antibody is evidenced by a colorimetric reaction.

Source: Mouse monoclonal

Species Reactivity: Human; others not tested

Clone: SN3b

Isotype: IgM/kappa

Total Protein Concentration: ~10 mg/ml. Call for lot specific Ig concentration.

Epitope/Antigen: CD24

Cellular Localization: Cytoplasmic and/or cell membrane

Positive Control: Some breast cancers

Known Applications:

Immunohistochemistry (formalin-fixed paraffin-embedded tissues)

Supplied As: Buffer with protein carrier and preservative

Storage and Stability:

Store at 2°C to 8°C. Do not use after expiration date printed on vial. If reagents are stored under conditions other than those specified in the package insert, they must be verified by the user. Diluted reagents should be used promptly; any remaining reagent should be stored at 2°C to 8°C.

Protocol Recommendations:

Peroxide Block: Block for 5 minutes with Biocare's Peroxidized 1.

Pretreatment Solution (recommended): Diva or Reveal

Pretreatment Protocol:

Heat Retrieval Method:

Preheat the retrieval solution to 95°C for 30 minutes in Biocare's Decloaking Chamber. Then, place slides into the preheated solution and retrieve under pressure at 95°C for 40 minutes; alternatively, steam tissue sections for 45-60 minutes or use a water bath at 95°C for 40 minutes. Allow solution to cool for 20 minutes then wash in distilled water.

Protocol Recommendations Cont'd:

Protein Block (Optional): Incubate for 5-10 minutes at RT with Biocare's Background Punisher.

Primary Antibody: Incubate for 30 minutes at RT.

Probe: N/A

Polymer: Incubate for 30 minutes at RT with a secondary-conjugated polymer.

Chromogen: Incubate for 5 minutes at RT with Biocare's DAB - OR - Incubate for 5-7 minutes at RT with Biocare's Warp Red.

Counterstain: Counterstain with hematoxylin. Rinse with deionized water. Apply Tacha's Bluing Solution for 1 minute. Rinse with deionized water.

Technical Note:

This antibody has been standardized with Biocare's MACH 2 detection system. It can also be used on an automated staining system and with other Biocare polymer detection kits. Use TBS buffer for washing steps.

Limitations:

The optimum antibody dilution and protocols for a specific application can vary. These include, but are not limited to: fixation, heat-retrieval method, incubation times, tissue section thickness and detection kit used. Due to the superior sensitivity of these unique reagents, the recommended incubation times and titers listed are not applicable to other detection systems, as results may vary. The data sheet recommendations and protocols are based on exclusive use of Biocare products. Ultimately, it is the responsibility of the investigator to determine optimal conditions. The clinical interpretation of any positive or negative staining should be evaluated within the context of clinical presentation, morphology and other histopathological criteria by a qualified pathologist. The clinical interpretation of any positive or negative staining should be complemented by morphological studies using proper positive and negative internal and external controls as well as other diagnostic tests.

Quality Control:

Refer to CLSI Quality Standards for Design and Implementation of Immunohistochemistry Assays; Approved Guideline-Second edition (I/LA28-A2). CLSI Wayne, PA, USA (www.clsi.org). 2011

Precautions:

1. This antibody contains less than 0.1% sodium azide. Concentrations less than 0.1% are not reportable hazardous materials according to U.S. 29 CFR 1910.1200, OSHA Hazard communication and EC Directive 91/155/EC. Sodium azide (NaN₃) used as a preservative is toxic if ingested. Sodium azide may react with lead and copper plumbing to form highly explosive metal azides. Upon disposal, flush with large volumes of water to prevent azide build-up in plumbing. (Center for Disease Control, 1976, National Institute of Occupational Safety and Health, 1976) (4)
2. Specimens, before and after fixation, and all materials exposed to them should be handled as if capable of transmitting infection and disposed of with proper precautions. Never pipette reagents by mouth and avoid contacting the skin and mucous membranes with reagents and specimens. If reagents or specimens come in contact with sensitive areas, wash with copious amounts of water. (5)
3. Microbial contamination of reagents may result in an increase in nonspecific staining.
4. Incubation times or temperatures other than those specified may give erroneous results. The user must validate any such change.
5. Do not use reagent after the expiration date printed on the vial.
6. The MSDS is available upon request and is located at <http://biocare.net/support/msds/>.

CD24

Concentrated Monoclonal Antibody
Control Number: 901-323-011615

ISO
9001&13485
CERTIFIED

References:

1. Surowiak P, *et al.* CD24 expression is specific for tamoxifen-resistant ductal breast cancer cases. *Anticancer Res.* 2006 Jan-Feb; 26(1B):629-34.
2. Baumann P, *et al.* CD24 expression causes the acquisition of multiple cellular properties associated with tumor growth and metastasis. *Cancer Res.* 2005 Dec 1; 65(23):10783-93.
3. Dontu G, Liu S, Wicha MS. Stem cells in mammary development and carcinogenesis: implications for prevention and treatment. *Stem Cell Rev.* 2005; 1(3):207-13.
4. Center for Disease Control Manual. Guide: Safety Management, NO. CDC-22, Atlanta, GA. April 30, 1976 "Decontamination of Laboratory Sink Drains to Remove Azide Salts."
5. Clinical and Laboratory Standards Institute (CLSI). Protection of Laboratory workers from occupationally Acquired Infections; Approved guideline-Third Edition CLSI document M29-A3 Wayne, PA 2005.

Troubleshooting:

Follow the antibody specific protocol recommendations according to data sheet provided. If atypical results occur, contact Biocare's Technical Support at 1-800-542-2002.