

# Leukocyte Common Antigen (CD45RB)

Prediluted Monoclonal Antibody

Control Number: 901-091-102114

ISO  
9001&13485  
CERTIFIED

**Catalog Number:** PM 091 AA  
**Description:** 6.0 ml, prediluted  
**Dilution:** Ready-to-use  
**Diluent:** N/A

**Intended Use:**

For In Vitro Diagnostic Use

Leukocyte Common Antigen (CD45RB) [PD7/26] is a mouse monoclonal antibody that is intended for laboratory use in the qualitative identification of CD45RB protein by immunohistochemistry (IHC) in formalin-fixed paraffin-embedded (FFPE) human tissues. The clinical interpretation of any staining or its absence should be complemented by morphological studies using proper controls and should be evaluated within the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

**Summary and Explanation:**

CD45RB recognizes an antigen found on lymphoid cells. It reacts with B cells, T-cell subtypes, monocytes, macrophages, and is weakly expressed with granulocytes (2-3). The PD7/26 antibody was included in the 3rd International Workshop. Studies have shown CD45RB reacts with 3 of the 5 glycoproteins in the CD45 family (MW 190, 205 and 220 kDa). It is similar to CD45, except it is not expressed in Langerhan's cells and a small subset of T cells.

**Principle of Procedure:**

Antigen detection in tissues and cells is a multi-step immunohistochemical process. The initial step binds the primary antibody to its specific epitope. After labeling the antigen with a primary antibody, a secondary antibody is added to bind to the primary antibody. An enzyme label is then added to bind to the secondary antibody; this detection of the bound antibody is evidenced by a colorimetric reaction.

**Source:** Mouse monoclonal

**Species Reactivity:** Human; others not tested

**Clone:** PD7/26

**Isotype:** IgG1/kappa

**Total Protein Concentration:** ~10 mg/ml. Call for lot specific Ig concentration.

**Epitope/Antigen:** CD45RB (Leukocyte Common Antigen)

**Cellular Localization:** Cell membrane

**Positive Tissue Control:** Tonsil or lymphoma

**Known Applications:**

Immunohistochemistry (formalin-fixed paraffin-embedded tissues)

**Supplied As:** Buffer with protein carrier and preservative

**Storage and Stability:**

Store at 2°C to 8°C. Do not use after expiration date printed on vial. If reagents are stored under conditions other than those specified in the package insert, they must be verified by the user.

**Protocol Recommendations:**

**Peroxide Block:** Block for 5 minutes with Biocare's Peroxidized 1.

**Pretreatment Solution (recommended):** Diva

**Pretreatment Protocol:**

Heat Retrieval Method:

Retrieve sections under pressure using Biocare's Decloaking Chamber, followed by a wash in distilled water; alternatively, steam tissue sections for 45-60 minutes. Allow solution to cool for 10 minutes then wash in distilled water.

**Protein Block (Optional):** Incubate for 5-10 minutes at RT with Biocare's Background Punisher.

**Primary Antibody:** Incubate for 30 minutes at RT.

**Probe:** Incubate for 10 minutes at RT with a secondary probe.

**Protocol Recommendations Cont'd:**

**Polymer:** Incubate for 10 minutes at RT with a tertiary polymer.

**Chromogen:** Incubate for 5 minutes at RT with Biocare's DAB - OR - Incubate for 5-7 minutes at RT with Biocare's Warp Red.

**Counterstain:** Counterstain with hematoxylin. Rinse with deionized water. Apply Tacha's Bluing Solution for 1 minute. Rinse with deionized water.

**Technical Note:**

This antibody has been standardized with Biocare's MACH 4 detection system. It can also be used on an automated staining system and with other Biocare polymer detection kits. Use TBS buffer for washing steps.

**Limitations:**

The optimum antibody dilution and protocols for a specific application can vary. These include, but are not limited to: fixation, heat-retrieval method, incubation times, tissue section thickness and detection kit used. Due to the superior sensitivity of these unique reagents, the recommended incubation times and titers listed are not applicable to other detection systems, as results may vary. The data sheet recommendations and protocols are based on exclusive use of Biocare products. Ultimately, it is the responsibility of the investigator to determine optimal conditions. The clinical interpretation of any positive or negative staining should be evaluated within the context of clinical presentation, morphology and other histopathological criteria by a qualified pathologist. The clinical interpretation of any positive or negative staining should be complemented by morphological studies using proper positive and negative internal and external controls as well as other diagnostic tests.

**Quality Control:**

Refer to CLSI Quality Standards for Design and Implementation of Immunohistochemistry Assays; Approved Guideline-Second edition (I/LA28-A2) CLSI Wayne, PA USA (www.clsi.org). 2011.

**Precautions:**

1. This antibody contains less than 0.1% sodium azide. Concentrations less than 0.1% are not reportable hazardous materials according to U.S. 29 CFR 1910.1200, OSHA Hazard communication and EC Directive 91/155/EC. Sodium azide (NaN<sub>3</sub>) used as a preservative is toxic if ingested. Sodium azide may react with lead and copper plumbing to form highly explosive metal azides. Upon disposal, flush with large volumes of water to prevent azide build-up in plumbing. (Center for Disease Control, 1976, National Institute of Occupational Safety and Health, 1976) (5)
2. Specimens, before and after fixation, and all materials exposed to them should be handled as if capable of transmitting infection and disposed of with proper precautions. Never pipette reagents by mouth and avoid contacting the skin and mucous membranes with reagents and specimens. If reagents or specimens come in contact with sensitive areas, wash with copious amounts of water. (6)
3. Microbial contamination of reagents may result in an increase in nonspecific staining.
4. Incubation times or temperatures other than those specified may give erroneous results. The user must validate any such change.
5. Do not use reagent after the expiration date printed on the vial.
6. The MSDS is available upon request and is located at <http://biocare.net/support/msds/>.

**Troubleshooting:**

Follow the antibody specific protocol recommendations according to data sheet provided. If atypical results occur, contact Biocare's Technical Support at 1-800-542-2002.

## Leukocyte Common Antigen (CD45RB)

Prediluted Monoclonal Antibody

Control Number: 901-091-102114

ISO  
9001&13485  
CERTIFIED**References:**

1. Basadonna GP, *et al.* Antibody-mediated targeting of CD45 isoforms: a novel immunotherapeutic strategy. *Proc Natl Acad Sci U S A.* 1998 Mar 31;95(7):3821-6.
2. Hart IJ, *et al.* Subpopulations of guinea-pig T lymphocytes defined by isoforms of the leucocyte common antigen. *Immunology.* 1992 Nov;77(3):377-84.
3. Kurtin PJ, Pinkus GS. Leukocyte common antigen—a diagnostic discriminant between hematopoietic and nonhematopoietic neoplasms in paraffin sections using monoclonal antibodies: correlation with immunologic studies and ultrastructural localization. *Hum Pathol.* 1985 Apr;16(4):353-65.
4. Warnke RA, *et al.* Diagnosis of human lymphoma with monoclonal anti-leukocyte antibodies. *N Eng J Med.* 1983 Nov 24;309(21):1275-81.
5. Center for Disease Control Manual. Guide: Safety Management, NO. CDC-22, Atlanta, GA. April 30, 1976 "Decontamination of Laboratory Sink Drains to Remove Azide Salts."
3. Clinical and Laboratory Standards Institute (CLSI). Protection of Laboratory workers from occupationally Acquired Infections; Approved guideline-Third Edition CLSI document M29-A3 Wayne, PA 2005.