

## COX-2

Concentrated and Prediluted Rabbit Monoclonal Antibody  
902-306-030618

**BIOCARE**  
M E D I C A L

<b>Catalog Number:</b>	<b>ACR 306 A</b>	<b>APR 306 AA</b>
<b>Description:</b>	0.1 ml, concentrated	6.0 ml prediluted
<b>Dilution:</b>	1:50	Ready-to-use
<b>Diluent:</b>	Da Vinci Green	N/A

### Intended Use:

For Research Use Only. Not for use in diagnostic procedures.

### Summary and Explanation:

Cyclooxygenase (COX-2) is an inducible enzyme involved in production of prostaglandins in inflammatory processes. It is involved in the response of cells to growth factors, tumor promoters, and cytokines that induce its expression. Given its role in synthesizing prostaglandins, COX-2 is therefore of interest in studying immune response regulation. COX-2 is induced by a wide variety of stimuli and was initially identified as immediate-early growth response gene. There is now increasing evidence that a constitutive expression of COX-2 plays a role in development and progression of malignant epithelial tumors. In studies, COX-2 positivity retained independent roles in predicting a poor chance of response to treatment. COX-2 positive patients had a shorter overall survival rate than COX-2 negative patients.

### Principle of Procedure:

Antigen detection in tissues and cells is a multi-step immunohistochemical process. The initial step binds the primary antibody to its specific epitope. After labeling the antigen with a primary antibody, an enzyme labeled polymer is added to bind to the primary antibody. This detection of the bound antibody is evidenced by a colorimetric reaction.

**Source:** Rabbit monoclonal

**Species Reactivity:** Human, mouse and rat

**Clone:** SP21

**Isotype:** IgG

**Total Protein Concentration:** ~10 mg/ml. Lot specific Ig concentration is not available.

**Epitope/Antigen:** Cyclooxygenase-2

**Cellular Localization:** Cytoplasmic

**Positive Tissue Control:** Breast, colon or lung carcinoma

### Known Applications:

Immunohistochemistry (formalin-fixed paraffin-embedded tissues)

**Supplied As:** Buffer with protein carrier and preservative

### Storage and Stability:

Store at 2°C to 8°C. Do not use after expiration date printed on vial. If reagents are stored under conditions other than those specified in the package insert, they must be verified by the user. Diluted reagents should be used promptly; any remaining reagent should be stored at 2°C to 8°C.

### Staining Protocol Recommendations:

**Peroxide Block:** Block for 5 minutes with Biocare's Peroxidized 1.

**Pretreatment:** Perform heat retrieval using Biocare's Diva or Reveal Decloaker. Refer to the Diva or Reveal Decloaker product data sheet for specific instructions.

**Protein Block (Optional):** Incubate for 5-10 minutes at RT with Biocare's Background Punisher.

**Primary Antibody:** Incubate for 30 minutes at RT.

**Probe:** N/A

**Polymer:** Incubate for 30 minutes at RT with a secondary-conjugated polymer.

**Chromogen:** Incubate for 5 minutes at RT with Biocare's DAB - OR - Incubate for 5-7 minutes at RT with Biocare's Warp Red.

### Counterstain:

Counterstain with hematoxylin. Rinse with deionized water. Apply Tacha's Bluing Solution for 1 minute. Rinse with deionized water.

### Technical Note:

This antibody has been standardized with Biocare's MACH 2 detection system. Use TBS buffer for washing steps.

### Limitations:

This product is provided for Research Use Only (RUO) and is not for use in diagnostic procedures. Suitability for specific applications may vary and it is the responsibility of the end user to determine the appropriate application for its use.

### Precautions:

1. This antibody contains less than 0.1% sodium azide. Concentrations less than 0.1% are not reportable hazardous material according to U.S. 29 CFR 1910.1200, OSHA Hazard communication and EC Directive 91/155/EC. Sodium azide (NaN<sub>3</sub>) used as a preservative is toxic if ingested. Sodium azide may react with lead and copper plumbing to form highly explosive metal azides. Upon disposal, flush with large volumes of water to prevent azide build-up in plumbing. (Center for Disease Control, 1976, National Institute of Occupational Safety and Health, 1976) (4)

2. Specimens, before and after fixation, and all materials exposed to them should be handled as if capable of transmitting infection and disposed of with proper precautions. Never pipette reagents by mouth and avoid contacting the skin and mucous membranes with reagents and specimens. If reagents or specimens come in contact with sensitive areas, wash with copious amounts of water. (5)

3. Microbial contamination of reagents may result in an increase in nonspecific staining.

4. Incubation times or temperatures other than those specified may give erroneous results. The user must validate any such change.

5. Do not use reagent after expiration date printed on the vial.

6. The SDS is available upon request and is located at <http://biocare.net>.

### Technical Support:

Contact Biocare's Technical Support at 1-800-542-2002 for questions regarding this product.

### References:

1. Boland GP, *et al.* COX-2 expression is associated with an aggressive phenotype in ductal carcinoma in situ. *Br J Cancer.* 2004 Jan 26; 90(2):423-9.

2. Laga AC, Zander DS, Cagle PT. Prognostic significance of cyclooxygenase 2 expression in 259 cases of non-small cell lung cancer. *Arch Pathol Lab Med.* 2005 Sep; 129(9):1113-7.

3. Soumaoro LT, *et al.* Cyclooxygenase-2 expression: a significant prognostic indicator for patients with colorectal cancer. *Clin Cancer Res.* 2004 Dec 15; 10(24):8465-71.

4. Center for Disease Control Manual. Guide: Safety Management, NO. CDC-22, Atlanta, GA. April 30, 1976 "Decontamination of Laboratory Sink Drains to Remove Azide Salts."

5. Clinical and Laboratory Standards Institute (CLSI). Protection of Laboratory Workers from Occupationally Acquired Infections; Approved Guideline-Fourth Edition CLSI document M29-A4 Wayne, PA 2014.



60 Berry Drive  
Pacheco, CA 94553  
USA

Rev: 062117

Tel: 800-799-9499 | [www.biocare.net](http://www.biocare.net) | Fax: 925-603-8080