

ROS1 [EPMGHR2]

Concentrated and Prediluted Rabbit Monoclonal Antibody
902-3240-060321

BIOCARE
M E D I C A L

Available Product Formats				
Format	Catalog Number	Description	Dilution	Diluent
Concentrate	ACR 3240 A, B	0.1, 0.5 mL	1:100	Renoir Red
Predilute	APR 3240 AA	6.0 mL	Ready-to-use	N/A
Q Series— For Leica BOND-III	ALR 3240 G7	7.0 mL	Ready-to-use	N/A

Intended Use:

For Research Use Only. Not for use in diagnostic procedures.

Summary and Explanation:

The *ROS1* gene encodes ROS proto-oncogene 1, a receptor tyrosine kinase. The ROS1 protein is an enzyme with a mass of 263.9 kDa. *ROS1* gene rearrangements have been implicated in multiple cancer indications including non-small cell lung cancer (NSCLC), gastric cancer, glioblastoma multiforme, ovarian cancer, and colorectal cancer (1). ROS1 protein fusion expression leads to oncogenic transformation, most likely through constitutive activation of the tyrosine kinase domain (2). *ROS1* gene rearrangements occur in 1-2% of NSCLC, and clinical trial data indicates that tumors positive for ROS1 protein fusions are sensitive to kinase inhibitors such as crizotinib, and entrectinib (3,4). For detection of positive ROS1 fusions, IHC testing has been shown 100% sensitivity and 92-94% specificity in recent studies, proving its efficacy as a screening tool for ROS1-targeted therapies (5,6).

Principle of Procedure:

Antigen detection in tissues and cells is a multi-step immunohistochemical process. The initial step binds the primary antibody to its specific epitope. After labeling the antigen with a primary antibody, a one-step or two-step detection procedure can be applied. A one-step procedure will feature an enzyme labeled polymer that binds the primary antibody. A two-step procedure will feature a linker antibody added to bind to the primary antibody. An enzyme-labeled polymer is then added to bind the linker antibody. These detections of the bound antibodies are evidenced by a colorimetric reaction.

Source: Rabbit monoclonal

Species Reactivity: Human

Clone: EPMGHR2

Isotype: IgG

Protein Concentration: Call for lot specific Ig concentration.

Epitope/Antigen: Synthetic peptide within Human ROS1

Cellular Localization: Cytoplasmic

Positive Tissue Control: Lung cancer

Known Applications:

Immunohistochemistry (formalin-fixed paraffin-embedded tissues)

Supplied As: Buffer with protein carrier and preservative

Storage and Stability:

Store at 2°C to 8°C. The product is stable to the expiration date printed on the label, when stored under these conditions. Do not use after expiration date. Diluted reagents should be used promptly; any remaining reagent should be stored at 2°C to 8°C.

Staining Protocol Recommendations (intelliPATH FLX® and manual use):

Peroxide Block: Block for 5 minutes with Peroxidized 1.

Pretreatment: Perform heat retrieval using Diva Decloaker. Refer to the Diva Decloaker product data sheet for specific instructions.

Protein Block (Optional): Incubate for 5-10 minutes at RT with Background Punisher.

Primary Antibody: Incubate for 60 minutes at RT.

Probe: N/A

Polymer: Incubate for 30 minutes at RT with a secondary-conjugated polymer.

Chromogen: Incubate for 5 minutes at RT with Biocare's DAB - OR - Incubate for 5-7 minutes at RT with Warp Red.

Counterstain:

Counterstain with hematoxylin. Rinse with deionized water. Apply Tacha's Bluings Solution for 1 minute. Rinse with deionized water.

Staining Protocol Recommendations (intelliPATH FLX and manual use) Cont'd:

Technical Note:

This antibody, for intelliPATH FLX and manual use, has been standardized with MACH 4 detection system. Use TBS for washing steps.

Staining Protocol Recommendations (Q Series – For Leica BOND-III):

ALR3240 is intended for use with the Leica BOND-III. Refer to the User Manual for specific instructions for use. Recommended protocol parameters are as follows:

Protocol Name: IHC Protocol F

Detection: Bond Polymer Refine

HIER: 20 min with ER1

Peroxide Block: 5 min

Marker (Primary Antibody): 15 min

Post Primary: 8 min

Polymer: 8 min

Mixed DAB Refine: 10 min

Hematoxylin: 5 min

Limitations:

This product is provided for Research Use Only (RUO) and is not for use in diagnostic procedures. Suitability for specific applications may vary and it is the responsibility of the end user to determine the appropriate application for its use.

Precautions:

1. This antibody contains less than 0.1% sodium azide. Concentrations less than 0.1% are not reportable hazardous materials according to U.S. 29 CFR 1910.1200, OSHA Hazard communication and EC Directive 91/155/EC. Sodium azide (NaN₃) used as a preservative is toxic if ingested. Sodium azide may react with lead and copper plumbing to form highly explosive metal azides. Upon disposal, flush with large volumes of water to prevent azide build-up in plumbing. (Center for Disease Control, 1976, National Institute of Occupational Safety and Health, 1976) (7).

2. Specimens, before and after fixation, and all materials exposed to them should be handled as if capable of transmitting infection and disposed of with proper precautions. Never pipette reagents by mouth and avoid contacting the skin and mucous membranes with reagents and specimens. If reagents or specimens come into contact with sensitive areas, wash with copious amounts of water (8).

3. Microbial contamination of reagents may result in an increase in nonspecific staining.

4. Incubation times or temperatures other than those specified may give erroneous results. The user must validate any such change.

5. Do not use reagent after the expiration date printed on the vial.

6. The SDS is available upon request and is located at <http://biocare.net>.

Technical Support:

Contact Biocare's Technical Support at 1-800-542-2002 for questions regarding this product.

References:

1. Gainer JF, Shaw, AT. Novel Targets in Non-Small Cell Lung Cancer: ROS1 and RET Fusions. *Oncologist*. 2013;18(7):865-75.
2. Davies KD, Doebele RC. Molecular pathways: ROS1 fusion proteins in cancer. *Clin Cancer Res*. 2013 Aug 1;19(15):4040-5.



60 Berry Drive
Pacheco, CA 94553
USA

Rev. 062117

Tel: 800-799-9499 | www.biocare.net | Fax: 925-603-8080

ROS1 [EPMGHR2]

Concentrated and Prediluted Rabbit Monoclonal Antibody
902-3240-060321

BIOCARE
M E D I C A L

References Cont'd:

3. Bubendorf L, *et al.* Testing for ROS1 in non-small cell lung cancer: a review with recommendations. *Virchows Arch.* 2016 Nov;469(5):489-503.
4. Facchinetti F, Friboulet L. Profile of entrectinib and its potential in the treatment of ROS1-positive NSCLC: evidence to date. *Lung Cancer (Auckl).* 2019 Sep 9;10:87-94.
5. Sholl LM, *et al.* ROS1 immunohistochemistry for detection of ROS1-rearranged lung adenocarcinomas. *Am J Surg Pathol.* 2013 Sep;37(9):1441-9.
6. Shan L, *et al.* Detection of ROS1 gene rearrangement in lung adenocarcinoma: comparison of IHC, FISH and real-time RT-PCR. *PLoS One.* 2015 Mar 5;10(3):e0120422.
7. Center for Disease Control Manual. Guide: Safety Management, NO. CDC-22, Atlanta, GA. April 30, 1976 "Decontamination of Laboratory Sink Drains to Remove Azide Salts."
8. Clinical and Laboratory Standards Institute (CLSI). Protection of Laboratory Workers from Occupationally Acquired Infections; Approved Guideline-Fourth Edition CLSI document M29-A4 Wayne, PA 2014.

Q Series antibodies are developed solely by Biocare Medical LLC and do not imply approval or endorsement of Biocare antibodies by Leica Biosystems. Biocare and Leica Biosystems are not affiliated, associated or related in any way. Leica, Leica Biosystems, BOND-MAX and BOND-III are trademarks of Leica Biosystems.



60 Berry Drive
Pacheco, CA 94553
USA

Rev. 062117

Tel: 800-799-9499 | www.biocare.net | Fax: 925-603-8080