

MACH 2 Mouse HRP-Polymer

Micro-polymer detection
902-BRR520A-032323

BIOCARE
M E D I C A L

Available Product Formats	
Catalog Number	Volume
BRR520AG	25 mL
BRR520AH	25 mL
BRR520AL	100 mL
BRR520AMM	1L

Intended Use:

For Research Use Only. Not for use in diagnostic procedures.

Background Information:

The MACH 2 Mouse HRP-Polymer is designed using a one-step method for detecting mouse primary antibodies to form an antibody-enzyme complex. This complex is then visualized using an appropriate substrate/chromogen. In the one-step method a secondary antibody directly linked to the micro-polymer is applied.

Known Applications:

Immunohistochemistry (Formalin-fixed paraffin-embedded tissues)

Materials and Methods:

Reagents Provided:

Kit Catalog No.	Component Description	Quantity x Volume
BRR520AG	MACH 2 Mouse HRP-Polymer	1 x 6.0 mL
BRR520AH	MACH 2 Mouse HRP-Polymer	1 x 25 mL
BRR520AL	MACH 2 Mouse HRP-Polymer	1 x 100 mL
BRR520AMM	MACH 2 Mouse HRP-Polymer	1 x 1000 mL

Supplied As:

Buffered saline solution, pH 7.6-7.8, containing a protein carrier and less than 0.01% ProClin 300 and/or less than 0.5% ProClin 950 as a preservative. See Safety Data Sheet for additional details.

Reconstitution, Dilution and Mixing:

The micro-polymer detection kit reagent(s) are optimized and ready to use with Biocare antibodies and ancillary reagents. No reconstitution, mixing, dilution, or titration is required.

Species Reactivity:

Mouse IgG heavy and light chains.

Storage and Stability:

Store at 2°C to 8°C. The product is stable to the expiration date printed on the vial label when stored under these conditions. Do not use after expiration date. Storage under any condition other than those specified must be verified. The kit reagent(s) are ready-to-use and should not be diluted. The stability of user diluted reagent has not been established by Biocare.

Staining Protocol Recommendations:

The below are programming and protocol recommendations to assist the user when staining manually and/or using one of Biocare's Automated Staining Platforms for research applications. The user is responsible for further optimizations of the protocol.

1. Deparaffinization: Deparaffinize slides in Slide Brite or xylene. Hydrate slides in a series of graded alcohols to water.
2. Peroxide Block: Block for 5 minutes with Peroxidized 1.
3. Pretreatment Solution/Protocol: Please refer to the respective primary antibody data sheet for recommended pretreatment solution and protocol.

Staining Protocol Recommendations Cont'd:

4. Protein Block (Optional): Incubate for 10-15 minutes at room temperature (RT) with Background Sniper.
5. Primary Antibody: Please refer to the respective primary antibody data sheet for incubation time.
6. Polymer: Incubate for 30 minutes at RT with MACH 2 Mouse HRP-Polymer.
7. Chromogen: Incubate for 5 minutes at RT with Biocare's DAB.
8. Counterstain: Counterstain with hematoxylin. Rinse with deionized water. Apply Tacha's Bluing Solution for 1 minute. Rinse with deionized water.

Technical Note:

1. Use TBS for washing steps.


Limitations:

This product is provided for Research Use Only (RUO) and is not for use in diagnostic procedures. Suitability for specific applications may vary and it is the responsibility of the end user to determine the appropriate application for its use.

Precautions:


1. Kit reagents contain less than 0.05% ProClin 300 and/or less than 1% ProClin 950. Wear gloves and protective clothing and take reasonable precautions when handling as ProClin is classified as an irritant and may cause skin contact sensitization. Avoid contact with eyes, skin, and mucous membranes.
2. Handle materials of human or animal origin as potentially biohazardous and dispose of such materials with proper precautions. In the event of exposure, follow the health directives of the responsible authorities where used.^{2,3}
3. Specimens, before and after fixation, and all materials exposed to them should be handled as if capable of transmitting infection and disposed of with proper precautions. Never pipette reagents by mouth and avoid contacting the skin and mucous membranes with reagents and specimens. If reagents or specimens come into contact with sensitive areas, wash with copious amounts of water.⁴
4. Microbial contamination of reagents may result in an increase in nonspecific staining.
5. Incubation times or temperatures other than those specified may give erroneous results. The user must validate any such change.
6. Do not use reagent after the expiration date printed on the vial.
7. The reagent is optimized for use with Biocare antibodies and ancillary reagents. Refer to the primary antibody and other ancillary reagent instructions for use for recommended protocols and conditions for use.
8. Follow local and/or state authority requirements for method of disposal.
9. The SDS is available upon request and is located at <http://biocare.net>.

This micro-polymer detection kit contains components classified as indicated in the table below in accordance with Regulation (EC) No. 1272/2008.

Hazard	Code	Hazard Statement
	H317	May cause an allergic skin reaction.
N/A	H402 H412	Harmful to aquatic life. Harmful to aquatic life with long lasting effects.

Technical Support:

Contact Biocare's Technical Support at 1-800-542-2002 for questions regarding this product.

 Biocare Medical
60 Berry Drive
Pacheco, CA 94553
USA

TP v2 (01/31/2023)

Tel: 800-799-9499 | www.biocare.net | Fax: 925-603-8080

MACH 2 Mouse HRP-Polymer

Micro-polymer detection
902-BRR520A-032323

BIOCARE
M E D I C A L

References:

1. Occupational Safety and Health Standards: Occupational exposure to hazardous chemicals in laboratories. (29 CFR Part 1910.1450). Fed. Register.
2. Directive 2000/54/EC of the European Parliament and Council of 18 September 2000 on the protection of workers from risks related to exposure to biological agents at work.
3. Clinical and Laboratory Standards Institute (CLSI). Protection of Laboratory Workers from Occupationally Acquired Infections; Approved Guideline-Fourth Edition CLSI document M29-A4 Wayne, PA 2014.



Biocare Medical
60 Berry Drive
Pacheco, CA 94553
USA

TP v2 (01/31/2023)

Tel: 800-799-9499 | www.biocare.net | Fax: 925-603-8080